

New substd. furyl, thienyl or pyrrolyl carbonyl-guanidine derivs. - used
e.g. as cellular sodium proton exchange inhibitors, antiarrhythmic agents
and cell proliferation inhibitors

Patent Assignee: HOECHST AG (FARH)

Inventor: ALBUS U; KLEEMANN H; LANG J; SCHOLZ W; SCHWARK J; WEICHERT A;
LANG H; SCHWARK J R

Number of Countries: 028 Number of Patents: 017

Patent Family:

Patent No	Kind	Date	Applicat No	Kind	Date	Week
EP 676395	A2	19951011	EP 95105088	A	19950405	199546 B
DE 4412334	A1	19951019	DE 4412334	A	19940411	199547
AU 9516354	A	19951019	AU 9516354	A	19950407	199549
NO 9501405	A	19951012	NO 951405	A	19950410	199549
FI 9501681	A	19951012	FI 951681	A	19950407	199601
JP 7291927	A	19951107	JP 95107811	A	19950410	199602
CA 2146707	A	19951012	CA 2146707	A	19950410	199607
ZA 9502930	A	19960327	ZA 952930	A	19950410	199619
EP 676395	A3	19960306	EP 95105088	A	19950405	199624
HU 71616	T	19960129	HU 951035	A	19950410	199738
CN 1117044	A	19960221	CN 95104391	A	19950410	199742
NZ 270894	A	19970922	NZ 270894	A	19950407	199745
AU 683722	B	19971120	AU 9516354	A	19950407	199804
US 5698581	A	19971216	US 95418434	A	19950407	199805
NO 304426	B1	19981214	NO 951405	A	19950410	199905
TW 349941	A	19990111	TW 95102001	A	19950303	199923
IL 113310	A	20000629	IL 113310	A	19950410	200047

Priority Applications (No Type Date): DE 4412334 A 19940411

Cited Patents: No-SR.Pub; 3.Jnl.Ref; DE 1965267; DE 2055727; EP 416499; EP
556672; EP 556673; EP 556674; EP 577024; EP 589336; EP 590455; EP 622356;
EP 639573; JP 44030268; WO 9304048

Patent Details:

Patent No Kind Lan Pg Main IPC Filing Notes

EP 676395 A2 G 48 C07D-207/40

Designated States (Regional): AT BE CH DE DK ES FR GB GR IE IT LI LU NL
PT SE

DE 4412334 A1 41 C07D-207/416

AU 9516354 A C07D-207/416

NO 9501405 A C07D-207/416

FI 9501681 A C07D-207/34

JP 7291927 A 28 C07D-207/416

CA 2146707 A C07D-207/34

ZA 9502930	A	70 C07D-000/00
EP 676395	A3	C07D-207/40
HU 71616	T	C07D-207/416
CN 1117044	A	A61K-031/44
NZ 270894	A	C07D-207/34
AU 683722	B	C07D-207/416 Previous Publ. patent AU 9516354
US 5698581	A	22 A61K-031/38
NO 304426	B1	C07D-207/337 Previous Publ. patent NO 9501405
TW 349941	A	C07D-207/34
IL 113310	A	C07D-207/34

Abstract (Basic): EP 676395 A

Heteroaryl-guanidine derivs. of formula (I) and their salts are new. A = S(O)m, O or NR5; m = 0, 1 or 2; R5 = H, 1-8C alkyl or CmH2mR81; R81 = 3-8C cycloalkyl, phenyl (opt. substd. by 1-3 of F, Cl, CF3, Me, OMe, and NR82R83) or 1-9C heteroaryl (bonded via C or N and opt. substd. by 1-3 of F, Cl, CF3, Me, OMe, OH, NH2, NHMe and NMe2); R82, R83 = H or Me; one of R1, R2 = CO-N=C(NH2)2; the other = H, F, Cl, Br, I, 1-3C alkyl, OR6, 1-4C perfluoroalkyl, CO-N=C(NH2)2 or NR6R7; R6, R7 = H or 1-3C alkyl; R3, R4 = (i) H, F, Cl, Br, I, CN, X(CH2)m(1-6C) perfluoroalkyl, X(CH2)mF, S(O)mR8, CONR9R10, COR11, SO2NR12R13; (ii) 1-8C alkyl, CmH2mR81; (iii) 1-9C heteroaryl (opt. substd. by 1-3 of F, Cl, CF3, Me, OMe, OH, NH2, NHMe and NMe2); (iv) -Y-C6H4-(CO)i-(CHOH)j-(CHOH)k-R23; (v) H, F, Cl, Br, I, CN, 1-8C alkyl, 1-8C perfluoroalkyl, 3-8C alkenyl, CgH2g-R26; SR29, OR30, NR31R32, CR33R34R35; (vii) -W-C6H4-R97; (viii) S(O)mR37, SO2NR38R39; (ix) X1R46; (x) SR64, OR65, NHR66, NR67R68, CHRR69R70, CR54R55-OH, Ctriple bondC-R56 C(R58) C-R57 (sic), (CR59R60)u-CO-(CR61R62)v-R63; (xi) SO2NHR76; or (xii) NR84R85; X = O, S or NR14; R14 = H or 1-3C alkyl; R8 = 1-5C alkyl, 3-6C alkenyl, CnH2nR15 or CF3; R9, R11, R12 = H or as R8; n = 0-4; R15 = 3-7C cycloalkyl or phenyl (opt. substd. by 1-3 of F, Cl, CF3, Me, OMe and NR16R17); R16, R17 = H or 1-4C alkyl; R10, R13 = H or 1-4C alkyl; or R9+R10 or R12+R13 = (CH2)4 or (CH2)5 in which one CH2 may be replaced by O, S, NH, NMe or N-benzyl; R18 = 3-8C cycloalkyl or phenyl (opt. substd. by 1-3 of F, Cl, CF3, Me, OMe and NR19R20); R19, R20 = H or Me; Y = O, S or NR22; h = 0 or 1; i, j, k = 0-4; provided that h, i and k are not all 0; R22, R23 = H or 1-3C alkyl; g = 0-4; R26 = 3-8C cycloalkyl, phenyl, biphenyl, or naphthyl (where aromatics are opt. substd. by 1-3 of F, Cl, CF3, Me, OMe and NR27R28); R27, R28 = H, 1-4C alkyl or 1-4C perfluoroalkyl; R29-R31, R33 = -(CH2)m-(1-9C) heteroaryl (opt. substd. as in R81); R32, R34, R35 = H, 1-4C alkyl, 1-4C perfluoroalkyl, or as R29; R96 = heteroaryl as defined for R81, or benzyl; W = O, S or NR36; R36 = H or 1-4C alkyl; R37 = 1-8C alkyl, 1-8C perfluoroalkyl, 3-8C alkenyl or -CsH2s-R40; s = 0-4; R40 = as R26; R38 = H, 1-8C alkyl, 1-8C perfluoroalkyl, 3-8C alkenyl or

-CwH₂w-R₂₆; R₃₉ = H, 1-4C alkyl or 1-4C perfluoroalkyl; or R₃₈+R₃₉ = (CH₂)₄ or (CH₂)₅, in which one CH₂ may be replaced by O, S, NH, NMe or N-benzyl; X₁ = O, S, NR₄₇, (D=O)A'- or NR₄₈C=MN*(R₄₉)-; M = O or S; A' = O or NR₅₀; D = C or SO; R₄₆, R₄₉ = 1-8C alkyl, 3-8C alkenyl, -(CH₂)_b-(1-7C)perfluoroalkyl or -C_xH_{2x}-R₂₆; b = 0 or 1; x = 0-4; R₄₇, R₄₈, R₅₀ = H, 1-4C alkyl or 1-4C perfluoroalkyl; or R₄₆+R₄₇ or R₄₆+R₄₈ = (CH₂)₄ or (CH₂)₅ in which CH₂ may be replaced by O, S, NH, NMe or N-benzyl; A' and N* are bonded to the phenyl ring of the benzoylguanidine structure; R₆₄-R₆₇, R₆₉ = -(CH₂)_y-(CHOH)_z-(CH₂)_q-(C_uH_{2u}O)_t-R₇₁ or -(CH₂)_{b'}-O-(CH₂CH₂O)_{c'}-R₇₂; R₇₁, R₇₂ = H or Me; b', c' are not defined; u, t = 1-4; v, y, z, a' = 0-4; R₆₈, R₇₀, R₅₄, R₅₅ = H or 1-6C alkyl; or CR₆₉R₇₀ or CR₅₄R₅₅ = 3-8C cycloalkylidene; R₆₃ = H, 1-6C alkyl, 3-8C cycloalkyl or -CeH_{2e}-R₇₃; e = 0-4; R₈₀ = 5-7C cycloalkyl or phenyl (opt. substd. by 1-3 of F, Cl, CF₃, OMe and 1-4C alkyl); or R₇₇+R₇₈ = (CH₂)₄ or (CH₂)₅, in which one CH₂ may be replaced by O, S, NH, NMe or N-benzyl; R₇₉ = as R₇₇; or amidino; R₈₄, R₈₅ = H or 1-4C alkyl; or R₈₄+R₈₅ = (CH₂)₄ or (CH₂)₅ in which one CH₂ may be replaced by O, S, NH, NMe or N-benzyl or 1 or 2 CH₂ gps. may be replaced by CH-Cd'H_{2d'}+1; d' is not defined. Cpds. (I; A = O; R₁ = -CON=C(NH₂)₂; R₂, R₃ = H; R₄ = H, Me or Et) are excluded.

USE - (I) are used for treatment of arrhythmia or shock states; for treatment or prophylaxis of cardiac infarct, angina pectoris, cardiac ischaemia, ischaemic states of the peripheral and central nervous system, stroke or ischaemic states of the peripheral organs and limbs; and adjuvant during surgical operations and organ transplants; in preservation and storage of transplants; for treatment of diseases in which cell proliferation is a prim. or sec. cause, esp. atherosclerosis, complications following diabetes, cancer, fibrotic diseases, (e.g. fibrosis of the lungs, liver or kidneys) or prostatic hyperplasia; and as reagents for inhibiting Na⁺/H⁺ exchange and for diagnosis of hypertension and proliferative diseases (all claimed). More generally (I) inhibit the cellular Na⁺/H⁺ exchange mechanism and cell proliferation and are useful for combatting oxygen deficiency states, pathological hypoxia and ischaemia. They are esp. useful as antiarrhythmic agents.

Daily dose is 0.001-10 (pref. 0.01-1) mg orally, parenterally, rectally or by inhalation.

ADVANTAGE - (I) have good antiarrhythmic activity, without undesirable salidiuretic side effects, potent cellular Na⁺/H⁺ exchange inhibiting activity and good water solubility (facilitating i.v. admin.).

(Dwg. 0/0)

Segment: CPI